



## Observations of the boundary-layer height and the wind profile in the marine boundary layer

Floors, Rogier Ralph; Gryning, Sven-Erik; Peña, Alfredo

*Published in:*  
EMS Annual Meeting Abstracts

*Publication date:*  
2015

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Floors, R. R., Gryning, S-E., & Peña, A. (2015). Observations of the boundary-layer height and the wind profile in the marine boundary layer. In *EMS Annual Meeting Abstracts* (Vol. 11). [EMS2014-500]

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## **Observations of the boundary-layer height and the wind profile in the marine boundary layer**

Rogier Floors, Sven-Erik Gryning, and Alfredo Peña

Department of Wind Energy, DTU, Roskilde, Denmark (rofl@dtu.dk)

Because of the large number of wind turbines that is planned to be installed offshore in the North Sea, there is an increasing interest in the representation of the wind profile in the marine boundary layer. We present an evaluation of the WRF model using a combination of mast measurements and remote sensing techniques. The wind, temperature and humidity and wave parameters are measured at the FINO 3 platform in the North Sea. The boundary layer height is estimated using a CL51 ceilometer from Vaisälä and the wind profile up to the PBL height is measured using a wind lidar from Leosphere from August 2013 onwards.

The backscatter coefficient was generally high and therefore the ceilometer was able to estimate the PBL height well. The PBL height showed a pronounced seasonal cycle: unstable conditions prevailed during autumn, while stable conditions were more common during spring. This caused a distinct behaviour of the wind profile, with relatively large wind shears during spring. The differences between the WRF model and the observations at FINO 3 will be discussed.